


[Web](#) [Images](#) [Video](#) [News](#) [Maps](#) [more »](#)

[Advanced Scholar Search](#)
[Scholar Preferences](#)
[Scholar Help](#)
Scholar All articles - **Recent articles** Results 1 - 10 of about 106 for **imide stationary phase enantiomer polysaccharide**. (0.09 seconds)

[Enantioselective chromatography as a powerful alternative for the preparation of drug enantiomers - all 5 versions »](#)

ER Francotte - Journal of Chromatography A, 2001 - Elsevier

... the **enantiomers** a tetracyclic **imide** derivative ... using the conventional mobile **phase** consisting of ... However, using the immobilized **stationary phase** with chloroform ...

[Cited by 62](#) - [Related Articles](#) - [Web Search](#)

[\[CITATION\] ... -arylpropionic acids, barbituric acids and benzodiazepines on chiralcel OJ-R chiral stationary phase](#)

A Van Overbeke, W Baeyens, H Oda, HY Aboul-Enen - Chromatographia, 1996 - Springer

... not be resolved into its **enantiomers** on either ... dipole-dipole stacking between the **imide** and ester di ... stereo- specific positioning on the **stationary phase** [11, 12 ...

[Cited by 12](#) - [Related Articles](#) - [Web Search](#) - [BL Direct](#)

[The use of polysaccharide phases in the separation of enantiomers. - all 2 versions »](#)

RW Stringham - Adv Chromatogr, 2006 - books.google.com

... simultaneous interactions with the **stationary phase** while the orientation of the other **enantiomer** greatly weakens ... extensive use of **polysaccharide** CSPs, these ...

[Cited by 5](#) - [Related Articles](#) - [Web Search](#) - [BL Direct](#)

[Facile synthesis and highly efficient resolution of a constrained cyclopropane analogue of ... - all 2 versions »](#)

Al Jiménez, P López, L Oliveros, C Cativiela - Tetrahedron, 2001 - Elsevier

... not necessary to isolate the intermediate **imide** due to ... The optical purity of the resolved **enantiomers** was assessed ... of the silica-bound **stationary phase** in the ...

[Cited by 10](#) - [Related Articles](#) - [Web Search](#) - [BL Direct](#)

[Screening of chiral stationary and mobile phases for method development of preparative separations](#)

E FRANCOTTE, D HUYNH, HA WETLI - GIT laboratory journal Europe, 2006 - cat.inist.fr

... **Imide**; **Imide**; Imida; Sédatif; Sedative; Sedante; Hypnotique ... Cromatografía HPLC; **Phase** stationnaire chirale; Chiral **stationary phase**; Fase estacionaria ...

[Web Search](#)

[... using urea-and imide-bearing chitosan phenylcarbamate derivatives as chiral stationary phases for ... - all 2 versions »](#)

C Yamamoto, M Fujisawa, M Kamigaito, Y Okamoto - CHIRALITY-NEW YORK-, 2008 - doi.wiley.com

... Preparation of **Stationary Phase** The packing materials were ... of racemates on carbamate-**imide** derivatives 2c ... the optical rotation of the first-eluted **enantiomer**. ...

[Related Articles](#) - [Web Search](#) - [BL Direct](#)

[Low-molecular-weight chiral cation exchangers: Novel chiral stationary phases and their application ... - all 6 versions »](#)

A Vienna - Electrophoresis, 2002 - doi.wiley.com

... like Pirkle-concept phases or **polysaccharide** phases are ... also advantageous in CEC **enantiomer** separation employing ... ie using an achiral **stationary phase** and the ...

[Cited by 23](#) - [Related Articles](#) - [Web Search](#) - [BL Direct](#)

[Synthesis and HPLC enantioseparation of the cyclopropane analogue of valine\(c 3 Val\) - all 3 versions »](#)

Al Jimenez, P Lopez, C Cativiela - Chirality, 2005 - doi.wiley.com

... and 1.0 g of the second-eluted **enantiomer** have been ... KEY WORDS: chiral HPLC; preparative HPLC resolution; chiral **stationary phase**; **polysaccharide**-derived **phase** ...

[Related Articles](#) - [Web Search](#)

[Effects from hydrogen bonding and rigidity on selectivity in tartaric acid-based chiral selectors ... - all 3 versions »](#)

J Oxelbark, IB Sellen - Chirality, 2003 - doi.wiley.com

... Development of new chiral **stationary phases** is thus of great ... **Imide** 2 was therefore proposed as an interesting candidate. ... Mobile **phase**: 5% 2-propanol in hexane. ...

[Cited by 1](#) - [Related Articles](#) - [Web Search](#) - [BL Direct](#)

[CITATION] Recent development in HPLC enantioseparation—a selected review

W Lindner - Chromatographia, 1987 - Springer

[Cited by 25](#) - [Related Articles](#) - [Web Search](#)

Key authors: [E Francotte](#) - [M Robson](#) - [M Cikalo](#) - [P Myers](#) - [M Euerby](#)

Google 

Result Page: [1](#) [2](#) [3](#) [4](#) [5](#) [6](#) [7](#) [8](#) [9](#) [10](#) [Next](#)

[Google Home](#) - [About Google](#) - [About Google Scholar](#)

©2008 Google